

Local Maximum Stable Sets Greedoids Stemmed from Very Well-Covered Graphs

Vadim E. Levit

Ariel University Center of Samaria, ISRAEL

E-mail: levitv@ariel.ac.il

Eugen Mandrescu

Holon Institute of Technology, ISRAEL

E-mail: eugen_m@hit.ac.il

Abstract

A *maximum stable set* in a graph G is a stable set of maximum cardinality. S is called a *local maximum stable set* of G , and we write $S \in \Psi(G)$, if S is a maximum stable set of the subgraph induced by the closed neighborhood of S . A greedoid $(V, \mathcal{F})$ is called a *local maximum stable set greedoid* if there exists a graph $G = (V, E)$ such that $\mathcal{F} = \Psi(G)$.

Nemhauser and Trotter Jr. [27], proved that any $S \in \Psi(G)$ is a subset of a maximum stable set of G . In [15] we have shown that the family $\Psi(T)$ of a forest T forms a greedoid on its vertex set. The cases where G is bipartite, triangle-free, well-covered, while $\Psi(G)$ is a greedoid, were analyzed in [17], [19], [21], respectively.

In this paper we demonstrate that if G is a very well-covered graph, then the family $\Psi(G)$ is a greedoid if and only if G has a unique perfect matching.

Keywords: very well-covered graph, local maximum stable set, perfect matching, greedoid, König-Egerváry graph.

1 Introduction

Throughout this paper $G = (V, E)$ is a simple (i.e., a finite, undirected, loopless and without multiple edges) graph with vertex set $V = V(G)$ and edge set $E = E(G)$. If $X \subset V$, then $G[X]$ is the subgraph of G spanned by X . If $A, B \subset V$ and $A \cap B = \emptyset$, then (A, B) stands for the set $\{e = ab : a \in A, b \in B, e \in E\}$. The *neighborhood* of a vertex $v \in V$ is the set $N(v) = \{u : u \in V \text{ and } vu \in E\}$. For $A \subset V$, we denote $N_G(A) = \{v \in V - A : N(v) \cap A \neq \emptyset\}$ and $N_G[A] = A \cup N(A)$, or shortly, $N(A)$ and $N[A]$. If $N(v) = \{u\}$, then v is a *pendant vertex* and uv is a *pendant edge* of G .

K_n, C_n, P_n denote respectively, the complete graph on $n \geq 1$ vertices, the chordless cycle on $n \geq 3$ vertices, and the chordless path on $n \geq 2$ vertices.

A *matching* in a graph $G = (V, E)$ is a set $M \subseteq E$ such that no two edges of M share a common vertex. A *maximum matching* is a matching of maximum cardinality. By $\mu(G)$ is denoted the size of a maximum matching. A matching is *perfect* if it saturates all the vertices of the graph.

If for every two incident edges of a cycle C exactly one of them belongs to a matching M , then C is called an M -alternating cycle [9]. It is clear that an M -alternating cycle should be of even length. A matching M in G is called *alternating cycle-free* if G has no M -alternating cycle. Alternating cycle-free matchings for bipartite graphs were first defined in [9]. For example, the matching $\{ab, cd, ef\}$ of the graph G from Figure 1 is alternating cycle-free.




Figure 1: The unique cycle of H is alternating with respect to the matching $\{yv, tx\}$.

A matching $M = \{a_i b_i : a_i, b_i \in V(G), 1 \leq i \leq k\}$ of graph G is called a *uniquely restricted matching* if M is the unique perfect matching of $G[\{a_i, b_i : 1 \leq i \leq k\}]$ [7].

Theorem 1.1 [7] *A matching M in a graph G is uniquely restricted if and only if G contains no alternating cycle with respect to M , i.e., M is alternating cycle-free.*

For instance, all the maximum matchings of the graph G in Figure 1 are uniquely restricted, while the graph H from the same figure has both uniquely restricted maximum matchings (e.g., $\{uv, xw\}$) and non-uniquely restricted maximum matchings (e.g., $\{xy, tv\}$).

A *stable set* in G is a set of pairwise non-adjacent vertices. A stable set of maximum size will be referred to as a *maximum stable set* of G , and the *stability number* of G , denoted by $\alpha(G)$, is the cardinality of a maximum stable set in G . Let $\Omega(G)$ stand for the set of all maximum stable sets of G .

In general, $\alpha(G) \leq \alpha(G - e)$ and $\mu(G - e) \leq \mu(G)$ holds for any edge e of a graph G . An edge e of G is α -critical (μ -critical) if $\alpha(G) < \alpha(G - e)$ ($\mu(G) > \mu(G - e)$, respectively). It is worth observing that there is no general connection between the α -critical and μ -critical edges of a graph.

Recall that G is called a *König-Egerváry graph* provided $\alpha(G) + \mu(G) = |V(G)|$ [3], [32]. As a well-known example, every bipartite graph is a König-Egerváry graph [4], [11].

Theorem 1.2 *If G is a König-Egerváry graph, then the following assertions hold:*

- (i) [16] *every maximum matching is contained in $(S, V(G) - S)$, for each $S \in \Omega(G)$;*
- (ii) [18] *the α -critical edges are also μ -critical, and they coincide in a bipartite graph.*

A set $A \subseteq V(G)$ is a *local maximum stable set* of G if $A \in \Omega(G[N[A]])$ [15]; by $\Psi(G)$ we denote the set of all local maximum stable sets of the graph G .




Figure 2: $\{a\}, \{e, d\}, \{a, d, f\} \in \Psi(G)$, while $\{b\}, \{a, e\}, \{c, f\}$ are not in $\Psi(G)$.

The following theorem concerning maximum stable sets in general graphs, due to Nemhauser and Trotter Jr. [27], shows that for a special subgraph H of a graph G , some maximum stable set of H can be enlarged to a maximum stable set of G .

Theorem 1.3 [27] *Every local maximum stable set of a graph is a subset of a maximum stable set.*

Let us notice that the converse of Theorem 1.3 is not generally true. For instance, C_n has no proper local maximum stable set, for any $n \geq 4$. The graph G in Figure 2 shows another counterexample: any $S \in \Omega(G)$ contains some local maximum stable set, but these local maximum stable sets are of different cardinalities. As examples, $\{a, d, f\} \in \Omega(G)$ and $\{a\}, \{d, f\} \in \Psi(G)$, while for $\{b, e, g\} \in \Omega(G)$ only $\{e, g\} \in \Psi(G)$.

Definition 1.4 [1], [10] *A greedoid is a pair $(V, \mathcal{F})$, where $\mathcal{F} \subseteq 2^V$ is a non-empty set system satisfying the following conditions:*

- Accessibility: *for every non-empty $X \in \mathcal{F}$ there is an $x \in X$ such that $X - \{x\} \in \mathcal{F}$;*
- Exchange: *for $X, Y \in \mathcal{F}$, $|X| = |Y| + 1$, there is an $x \in X - Y$ such that $Y \cup \{x\} \in \mathcal{F}$.*

Definition 1.5 [23] *A greedoid $(V, \mathcal{F})$ is called a local maximum stable set greedoid if there exists a graph $G = (V, E)$ such that $\mathcal{F} = \Psi(G)$.*

In fact, the following theorem says that, in the case of local maximum stable set greedoids, it is enough to check only the accessibility property.

Theorem 1.6 [23] *If the family $\Psi(G)$ of a graph G satisfies the accessibility property, then $(V(G), \Psi(G))$ is a greedoid.*

In the sequel, we use $\mathcal{F}$ instead of $(V, \mathcal{F})$, as the ground set V will be, usually, the vertex set of some graph.

Theorem 1.7 [15] *The family of local maximum stable sets of a forest forms a greedoid on its vertex set.*

The conclusion of Theorem 1.7 is not specific for forests. For instance, the family $\Psi(G)$ of the graph G in Figure 3 is a greedoid.




Figure 3: Both G and H are bipartite, but only $\Psi(G)$ forms a greedoid.

Notice that $\Psi(H)$ is not a greedoid, where H is from Figure 3, because the accessibility property is not satisfied; e.g., $\{y, t\} \in \Psi(H)$, while $\{y\}, \{t\} \notin \Psi(H)$. In addition, one can see that all the maximum matchings of the graph G in Figure 3 are uniquely restricted, while the graph H from the same figure has both uniquely restricted maximum matchings (e.g., $\{uv, xw\}$) and non-uniquely restricted maximum matchings (e.g., $\{xy, tv\}$). It turns out that this is the reason that $\Psi(H)$ is not a greedoid, while $\Psi(G)$ is a greedoid.

Theorem 1.8 [17] *For a bipartite graph G , $\Psi(G)$ is a greedoid on its vertex set if and only if all its maximum matchings are uniquely restricted.*




Figure 4: $\Psi(G)$ is not a greedoid, $\Psi(H)$ is a greedoid.

The case of bipartite graphs owning a unique cycle, whose family of local maximum stable sets forms a greedoid is analyzed in [14].

The graphs from Figure 4 are non-bipartite König-Egerváry graphs, and all their maximum matchings are uniquely restricted. Let us remark that both graphs are also triangle-free, but only $\Psi(H)$ is a greedoid. It is clear that $\{b, c\} \in \Psi(G)$, while $G[N[\{b, c\}]]$ is not a König-Egerváry graph. As one can see from the following theorem, this observation is the real reason for $\Psi(G)$ not to be a greedoid.

Theorem 1.9 [19] *If G is a triangle-free graph, then $\Psi(G)$ is a greedoid if and only if all maximum matchings of G are uniquely restricted and the closed neighborhood of every local maximum stable set of G induces a König-Egerváry graph.*

Let X be a graph with $V(X) = \{v_i : 1 \leq i \leq n\}$, and $\{H_i : 1 \leq i \leq n\}$ be a family of graphs. Joining each $v_i \in V(X)$ to all the vertices of H_i , we obtain a new graph, called the *corona* of X and $\{H_i : 1 \leq i \leq n\}$ and denoted by $G = X \circ \{H_1, H_2, \dots, H_n\}$. For instance, see Figure 5. If $H_1 = H_2 = \dots = H_n = H$, we write $G = X \circ H$, and in this case, G is called the *corona* of X and H .




Figure 5: The corona $G = X \circ \{K_3, K_2, P_3, C_4\}$.

If each H_i is a complete graph, then $X \circ \{H_1, H_2, \dots, H_n\}$ is called the *clique corona* of X and $\{H_1, H_2, \dots, H_n\}$; notice that the clique corona is well-covered (and very well-covered, whenever $H_i = K_1, 1 \leq i \leq n$). Recall that G is *well-covered* if all its maximal stable sets have the same cardinality [28], and G is *very well-covered* if, in addition, it has no isolated vertices and $|V(G)| = 2\alpha(G)$ [5].

A number of classes of well-covered graphs were completely described (see, for instance, the following references: [5], [6], [8], [12], [29], [30], [31]).

Theorem 1.10 (i) [6] *Let G be a connected graph of girth ≥ 6 , which is isomorphic to neither C_7 nor K_1 . Then G is well-covered if and only if $G = H \circ K_1$, for some graph H of girth ≥ 6 .*

(ii) [2], [20] *Let G be a graph having girth ≥ 5 . Then G is very well-covered if and only if $G = H \circ K_1$, for some graph H of girth ≥ 5 .*

(iii) [12] *G is very well-covered if and only if G is a well-covered König-Egerváry graph.*

(iv) [33] *$G = X \circ \{H_1, H_2, \dots, H_n\}$ is well-covered if and only if all H_i are complete.*

It is easy to prove that every graph having a perfect matching consisting of pendant edges is very well-covered. The converse is not generally true (see, for instance, the graphs depicted in Figure 6). Moreover, there are well-covered graphs without perfect matchings; e.g., K_3 .



Figure 6: Very well-covered graphs with no perfect matching consisting of only pendant edges.

Theorem 1.11 [5] *For a graph G without isolated vertices the following are equivalent:*

- (i) G is very well-covered;
- (ii) there exists a perfect matching M in G that satisfies property P , i.e.,

“ $N(x) \cap N(y) = \emptyset$, and each $v \in N(x) - \{y\}$ is adjacent to all vertices of $N(y) - \{x\}$ ”

hold for every edge $xy \in M$;

- (iii) there exists at least one perfect matching in G and every perfect matching in G satisfies property P .

Various cases of well-covered graphs generating local maximum stable set greedoids, were treated in [21], [22], [24], [25].

Theorem 1.12 *Let $G = X \circ \{H_1, H_2, \dots, H_n\}$, where $H_1, H_2, \dots, H_n$ are non-empty graphs.*

- (i) [24] *if $G = P_n$ and all $H_i, 1 \leq i \leq n$, are complete graphs, then $\Psi(G)$ is a greedoid;*
- (ii) [21] *if $H_i = K_1, 1 \leq i \leq n$, then $\Psi(G)$ is a greedoid;*
- (iii) [22] *if all $H_1, H_2, \dots, H_n$ are complete graphs, then $\Psi(G)$ is a greedoid;*
- (iv) [25] *$\Psi(G)$ is a greedoid if and only if every $\Psi(H_i), i = 1, 2, \dots, n$, is a greedoid.*

It turns out that the property of having a unique maximum matching is of crucial importance for very-well covered graphs to generate local maximum stable set greedoids.

Theorem 1.13 [26] *Let G be a very well-covered graph of girth at least 4. Then $\Psi(G)$ is a greedoid if and only if G has a unique maximum matching.*

In this paper we completely characterize very well-covered graphs whose families of local maximum stable sets are greedoids.

2 Very well-covered graphs producing greedoids

Notice that $S_1 = \{a, b\}$ and $S_2 = \{c, d\}$ are stable sets in the graph G_1 from Figure 7, $S_1 \in \Psi(G_1)$, and both $G_1[N[S_1]]$ and $G_1[N[S_2]]$ are König-Egerváry graphs. On the other hand, $S_2 = \{x, y\} \in \Psi(G_2)$, where G_2 is from Figure 7, but $G[N[S_3]]$ is not a König-Egerváry graph.

Theorem 2.1 [20] *Let G be a very well-covered graph. Then $G[N[S]]$ is a König-Egerváry graph, for every $S \in \Psi(G)$.*




Figure 7: Non-bipartite well-covered graphs. Moreover, G_1 is very well-covered.

Concerning the graph G_1 from Figure 7, let us remark that $\{b, d\}, \{b, e\}$ are stable sets, $|\{b, d\}| < |N(\{b, d\})|$ and $|\{b, e\}| = |N(\{b, e\})|$, but only $\{b, e\} \in \Psi(G_1)$.

Lemma 2.2 *If S is a stable set in a very well-covered graph G , then $S \in \Psi(G)$ if and only if $|S| = |N(S)|$.*

Proof. According to Theorems 1.10(iii) and 1.11, G is a König-Egerváry graph having a perfect matching, say M . If S is a stable set in G , there must be some $A \in \Omega(G)$, such that $S \subseteq A$, because G is well-covered. By Theorem 1.2(i), we have that $M \subseteq (A, V(G) - A)$. Since M is a perfect matching, it follows that S is matched into $N(S)$, and further,

$$|S| = |M(S)| \leq |N(S)|,$$

where $M(S) = \{y \in V : xy \in M, x \in S\}$.

Let $S \in \Psi(G)$. According to Theorem 2.1, $G[N[S]]$ is a König-Egerváry graph, and consequently, we get that $|S| = |M(S)| \geq |N(S)|$. Hence, we infer that $|S| = |N(S)|$.

Conversely, let S be a stable set in G satisfying $|S| = |N(S)|$. Since S is matched by M into $N(S)$, we infer that the restriction of M to $G[N[S]]$ is a perfect matching. Therefore, $|S| = \alpha(G[N[S]])$, and this implies $S \in \Psi(G)$. ■




Figure 8: G_1 and G_2 are not very well-covered graphs. Only G_1 is well-covered.

Notice that the above lemma can fail in a non-very well-covered graph. For instance, $S = \{x, y\} \in \Psi(G_1)$, while $|S| < |N(S)|$, where G_1 is from Figure 8 and it is well-covered. Further, the sets $S_1 = \{a, c\}, S_2 = \{e, d\}$ and $S_3 = \{v, w\}$ belong to $\Psi(G_2)$, where G_2 is from Figure 8, and they satisfy:

$$|S_1| = |N(S_1)|, |S_2| < |N(S_2)|, \text{ and } |S_3| > |N(S_3)|.$$

Concerning the very well-covered graph G_1 from Figure 7, we see that $A, B \in \Psi(G_1)$, where $B = \{a\}, A = B \cup \{a\}$, and $|N(A)| = |N(B)| + 1$. The following lemma shows that in a very well-covered graph the existence of an accessibility chain is equivalent to the fact that one can have a chain of stable sets, where each additional vertex added to a stable set increases the size of its open neighborhood by exactly one element.

Lemma 2.3 *If $A = B \cup \{v\}$ is a stable set in a very well-covered graph G , and $B \in \Psi(G)$, then $A \in \Psi(G)$ if and only if $|N(A)| = |N(B)| + 1$.*

Proof. Assume that $A = B \cup \{v\} \in \Psi(G)$. Since G is very well-covered, by Lemma 2.2, it follows that $|N(A)| - |N(B)| = |A| - |B| = 1$.

Conversely, since

$$|N(A)| = |N(B)| + 1 = |B| + 1 = |A|,$$

Lemma 2.2 implies immediately that $A \in \Psi(G)$. ■

Lemma 2.3 fails for graphs that are not very well-covered, for instance, $B_1 = \{x, y\}$ and $A_1 = B_1 \cup \{v\}$ belong to $\Psi(G_1)$, but $|N(A_1)| = |N(B_1)| + 2$, where G_1 is from Figure 8, and also $B_2 = \{v\}$ and $A_2 = B_2 \cup \{w\}$ belong to $\Psi(G_2)$, but $|N(A_2)| = |N(B_2)|$, where G_2 is from Figure 8.

Let us notice that the graphs G_1, G_2 and G_3 from Figure 9 are very well-covered; by Theorem 1.8 or 1.9, neither $\Psi(G_2)$ nor $\Psi(G_3)$ is a greedoid. However, $\Psi(G_1)$ is a greedoid.



Figure 9: G_1, G_2, G_3 are very well-covered graphs. G_1 has a unique perfect matching.

Theorem 2.4 *Let G be a very well-covered graph. Then $\Psi(G)$ forms a greedoid if and only if G has a unique maximum matching.*

Proof. Suppose that $\Psi(G)$ forms a greedoid. By Theorem 1.11, G has at least one perfect matching, say M .

Since $\Psi(G)$ is a greedoid, every $S \in \Omega(G)$ has an accessibility chain

$$\{x_1\} \subset \{x_1, x_2\} \subset \dots \subset \{x_1, x_2, \dots, x_{\alpha-1}\} \subset \{x_1, x_2, \dots, x_{\alpha}\} = S.$$

Let us denote $S_i = \{x_1, x_2, \dots, x_i\}, 1 \leq i \leq \alpha$, and $S_0 = \emptyset$.

Since $S_{i-1} \in \Psi(G), S_i = S_{i-1} \cup \{x_i\} \in \Psi(G)$ and G is well-covered, Lemma 2.3 implies that $|N(x_i) - N[S_{i-1}]| = 1$, because

$$|N(x_i) - N[S_{i-1}]| = |N(S_i) - N(S_{i-1})| = |N(S_i)| - |N(S_{i-1})|.$$

Let $\{y_i\} = N(x_i) - N[S_{i-1}], 1 \leq i \leq \alpha$. Hence, $M = \{x_i y_i : 1 \leq i \leq \alpha\}$ is a maximum matching in G .

Let us validate that M is a uniquely restricted maximum matching in G .

We induct on $k = |S_k|$ in order to show that the restriction of M to $H_k = G[N[S_k]]$, which we denote by M_k , is a uniquely restricted maximum matching in H_k .

For $k = 1, S_1 = \{x_1\} \in \Psi(G)$ and this implies that $N(x_1) = \{y_1\}$, unless x_1 is an isolated vertex. In this case, $M_1 = \{x_1 y_1\}$ is a uniquely restricted maximum matching in H_1 . If x_1 is an isolated vertex, then $M_1 = \emptyset$ is a uniquely restricted maximum matching in H_1 .

Suppose that the assertion is true for all $j \leq k - 1$. Let us notice that

$$N[S_k] = N[S_{k-1}] \cup \{x_k\} \cup \{y_k\}.$$

As we know, $N(x_k) - N[S_{k-1}] = \{y_k\}$.

Since H_k is König-Egerváry graph, M_k is a maximum matching in H_k . The edge $x_k y_k$ is α -critical in H_k , because $\{y_k\} = N(x_k) - N[S_{k-1}]$, and hence, $x_k y_k$ is also μ -critical in H_k , according to Theorem 1.2(ii). Therefore, any maximum matching of H_k contains the edge $x_k y_k$. Since $M_k = M_{k-1} \cup \{x_k y_k\}$ and M_{k-1} is a uniquely restricted maximum matching in $H_{k-1} = H_k - \{x_k, y_k\}$, it follows that M_k is a uniquely restricted maximum matching in H_k .

Conversely, assume that G has a unique perfect matching, say M .

We show that $\Psi(G)$ satisfies the accessibility property, i.e., for every non-empty $X \in \Psi(G)$ there is an $x \in X$ such that $X - \{x\} \in \Psi(G)$.

Let $S \in \Psi(G)$. According to Theorem 1.3, there is some $A \in \Omega(G)$, such that $S \subseteq A$. By Theorem 1.2(i), $M \subseteq (A, V(G) - A)$ and this implies that S is matched into $N(S)$, and further, $|S| = |M(S)| \leq |N(S)|$, where $M(S) = \{y \in V : xy \in M, x \in S\}$. According to Theorem 2.1, $G[N[S]]$ is a König-Egerváry graph, and this fact ensures that $|S| = |M(S)| \geq |N(S)|$. Hence, we infer that $N(S) = M(S)$.

Suppose that S does not satisfies the accessibility property, i.e., $S - \{x\} \notin \Psi(G)$ for every $x \in S$. This implies that $N(S - \{x\}) = N(S)$, for every $x \in S$. Consequently, each vertex in $N(S)$ has at least two neighbors in S .

We show that there is an even cycle C in $G[N[S]]$, such that half of its edges are in M .

Let $x_1 y_1 \in M$ and $x_1 \in S$. Since $|N(y_1) \cap S| \geq 2$, there is a vertex, say x_2 , belonging to $N(y_1) \cap S$.

Let $x_2 y_2 \in M$; such an edge exists, because M matches S into $M(S)$. Now, since $|N(y_2) \cap S| \geq 2$, there is a vertex, say x_3 , belonging to $N(y_2) \cap S$. If $x_3 = x_1$, then the cycle C spanned by $\{x_1, y_1, x_2, y_2\}$ has half of its edges in M . If $x_3 \neq x_1$, then we consider the edge in M that saturates x_3 , say $x_3 y_3 \in M$. Since $G[N[S]]$ is finite, after a number of steps, we find some vertex in $N(S)$, say y_k , that is joined by an edge to some x_j for $j < k$. Clearly, the cycle C , with

$$V(C) = \{x_i, y_i : j \leq i \leq k\}$$

and

$$E(C) = \{x_i, y_i : j \leq i \leq k\} \cup \{y_i x_{i+1} : 1 \leq i \leq k-1\} \cup \{x_j y_k\}$$

is even and has half of its edges in M . Therefore, $M' = (M - E(C)) \cup (E(C) - M)$ is a perfect matching in G and $M \neq M'$, in contradiction with the uniqueness of M in G .

Consequently, $\Psi(G)$ satisfies the accessibility property, and, according to Theorem 1.6, $\Psi(G)$ is a greedoid. ■

Let us remark that the very well-covered graph G_1 in Figure 9 has a C_3 and a C_4 ; one edge of C_4 belongs to the unique perfect matching M of G_1 , but none of the edges of C_3 is included in M .

Lemma 2.5 *No edge of some C_q , for $q = 3$ or $q \geq 5$, belongs to a perfect matching in a very well-covered graph.*

Proof. If the graph G is very well-covered, then by Theorem 1.11, G has a perfect matching, say M , and each perfect matching satisfies Property P .

Let $xy \in M$. Then, Property P implies that $N(x) \cap N(y) = \emptyset$, i.e., xy belongs to no C_3 in G . Further, if $v \in N(x) - \{y\}$ and $u \in N(y) - \{x\}$, Property P assures that $vu \in E(G)$, i.e., xy belongs to no C_q , for $q \geq 5$. ■

The very well-covered graphs G_1 , G_2 , and G_3 from Figure 10 have chordless alternating cycles of length 4. In addition, G_3 has an alternating cycle of length 6, namely, $\{e_1, e_2, \dots, e_6\}$ is alternating with respect to the perfect matching $\{e_1, e_3, e_5\}$.




Figure 10: Very well-covered graphs, each having more than one perfect matching.

Lemma 2.6 *Let G be a very well-covered graph and M be one of its maximum matchings. There exists an alternating cycle with respect to M if and only if there is an alternating chordless cycle of length four with respect to M .*

Proof. According to Theorem 1.11, every maximum matching of G is perfect. Suppose C^1 is an alternating cycle with respect to a perfect matching $M = \{a_i b_i : 1 \leq i \leq |V(G)|/2\}$. Without loss of generality, assume that

$$C^1 = \{a_1, b_1, a_2, b_2, a_3, \dots, a_{k-1}, b_{k-1}, a_k, b_k\}$$

is a cycle on $2k > 4$ vertices with edges

$$E(C^1) = \{a_1 b_1, b_1 a_2, a_2 b_2, b_2 a_3, \dots, a_{k-1} b_{k-1}, b_{k-1} a_k, a_k b_k, b_k a_1\}.$$

Since $a_1 b_1 \in M$, $b_k \in N(a_1) - \{b_1\}$, $a_2 \in N(b_1) - \{a_1\}$, Property P implies that a_2 is adjacent to b_k . Thus the cycle on $2k - 2$ vertices $C^2 = \{a_2, b_2, a_3, \dots, a_{k-1}, b_{k-1}, a_k, b_k\}$ with edges

$$E(C^2) = \{a_2 b_2, b_2 a_3, \dots, a_{k-1} b_{k-1}, b_{k-1} a_k, a_k b_k, b_k a_2\}$$

is still alternating with respect to M . It is clear that reducing the size of the cycle in this way one can easily reach C^{k-1} of size $2k = 4$. According to Lemma 2.5, C^{k-1} is an induced cycle of length four.

The converse is evident. ■

The conclusion of Lemma 2.6 can be true for non-well-covered graphs; e.g., the perfect matching $\{e_1, e_2, e_3\}$ of the graph G_1 from Figure 11 admits alternating cycles of length six and chordless of length four. On the other hand, Lemma 2.6 can fail for well-covered graphs; e.g., the perfect matching $\{e_1, e_2, e_3\}$ of the graph G_2 from Figure 11 admits a unique alternating cycle of length six, while the perfect matching $\{e_1, e_2, e_3, e_4\}$ of the graph G_3 from Figure 11 admits an alternating cycle of length four that has chords.




Figure 11: G_1 is not well-covered. G_2, G_3 are well-covered, but not very well-covered graphs.

Theorem 2.7 *Let G be a very well-covered graph. Then the following are true:*

- (i) $\Psi(G)$ is a greedoid;
- (ii) G has a uniquely restricted maximum matching;
- (iii) G has an alternating cycle-free maximum matching;
- (iv) G has an alternating C_4 -free maximum matching;
- (v) every maximum matching in G is alternating cycle-free;
- (vi) every maximum matching in G is alternating C_4 -free;
- (vii) all maximum matchings of G are uniquely restricted.

Proof. Firstly, Theorem 1.11 implies that each maximum matching of G is perfect.

(i) $\implies$ (ii) Theorem 2.4 claims that G must have a unique perfect matching, say M . Clearly, M is a uniquely restricted maximum matching.

(ii) $\implies$ (iii) It is true, by Theorem 1.1.

(iii) $\implies$ (iv) Clear.

(iv) $\implies$ (v) In fact, G has a perfect matching, say M , which is alternating C_4 -free. Hence, by Lemma 2.6, M is alternating cycle-free. Consequently, by Theorem 1.1, G has no other maximum matchings, and thus the assertion (v) is true.

(v) $\implies$ (vi) Clear.

(vi) $\implies$ (vii) By Lemma 2.6 and Theorem 1.1, it follows that every maximum matching of G is uniquely restricted.

(vii) $\implies$ (i) Since all maximum matchings of G are both perfect and uniquely restricted, it follows that G has a unique perfect matching. Consequently, $\Psi(G)$ is a greedoid, according to Theorem 2.4. ■

Corollary 2.8 *Each very well-covered C_4 -free graph G has a unique maximum matching, and, consequently, produces a local maximum stable sets greedoid.*

Proof. Combining Theorem 1.11 and Lemma 2.6, we infer that G has a unique perfect matching. Hence, Theorem 2.4 ensures that $\Psi(G)$ is a greedoid. ■

3 Conclusions

In this paper we have proved that a very well-covered graph produces a local maximum stable set greedoid if and only if it has a unique perfect matching.




Figure 12: The well-covered graph G has $\{e_1, e_2, e_3, e_4, e_5\}$ as its unique perfect matching. G is not very well-covered, since $\alpha(G) < 5 = |V(G)|/2$.

Nevertheless, the assertion is not true for every well-covered graphs with a unique perfect matching; e.g., $\Psi(G)$ is not a greedoid, where G is the well-covered graph from Figure 12, because $\{x, y\} \in \Psi(G)$, while $\{x\}, \{y\} \notin \Psi(G)$. Theorem 1.12 points out to a number of examples of well-covered graphs whose families of local maximum stable graphs form greedoids. For general well-covered graphs we propose the following.

Problem 3.1 *Characterize well-covered graphs producing local maximum stable set greedoids.*

References

- [1] A. Björner, G. M. Ziegler, *Introduction to greedoids*, in N. White (ed.), *Matroid Applications*, 284-357, Cambridge University Press, 1992.
- [2] N. Dean, J. Zito, *Well-covered graphs and extendability*, Discrete Mathematics **126** (1994) 67-80.
- [3] R. W. Deming, *Independence numbers of graphs - an extension of the König-Egerváry theorem*, Discrete Mathematics **27** (1979) 23-33.
- [4] E. Egervary, *On combinatorial properties of matrices*, Mat. Lapok **38** (1931) 16-28.
- [5] O. Favaron, *Very well-covered graphs*, Discrete Mathematics **42** (1982) 177-187.
- [6] A. Finbow, B. Hartnell, R. J. Nowakowski, *A characterization of well-covered graphs of girth 5 or greater*, Journal of Combinatorial Theory, B **57** (1993) 44-68.
- [7] M. C. Golumbic, T. Hirst, M. Lewenstein, *Uniquely restricted matchings*, Algorithmica **31** (2001) 139-154.
- [8] B. Hartnell, M. D. Plummer, *On 4-connected claw-free well-covered graphs*, Discrete Applied Mathematics **64** (1996) 57-65.
- [9] S. Krogdahl, *The dependence graph for bases in matroids*, Discrete Mathematics **19** (1977) 47-59.
- [10] B. Korte, L. Lovász, R. Schrader, *Greedoids*, Springer-Verlag, Berlin, 1991.
- [11] D. König, *Graphen und Matrizen*, Mat. Lapok **38** (1931) 116-119.
- [12] V. E. Levit, E. Mandrescu, *Well-covered and König-Egerváry graphs*, Congressus Numerantium **130** (1998) 209-218.
- [13] V. E. Levit, E. Mandrescu, *On the structure of α -stable graphs*, Discrete Mathematics **236** (2001) 227-243.
- [14] V. E. Levit, E. Mandrescu, *Unicycle bipartite graphs with only uniquely restricted maximum matchings*, Proceedings of the Third International Conference on Combinatorics, Computability and Logic, (DMTCS'1), Springer, (C. S. Calude, M. J. Dinneen and S. Sburlan eds.) (2001) 151-158.
- [15] V. E. Levit, E. Mandrescu, *A new greedoid: the family of local maximum stable sets of a forest*, Discrete Applied Mathematics **124** (2002) 91-101.
- [16] V. E. Levit and E. Mandrescu, *On α^+ -stable König-Egerváry graphs*, Discrete Mathematics **263** (2003) 179-190.
- [17] V. E. Levit, E. Mandrescu, *Local maximum stable sets in bipartite graphs with uniquely restricted maximum matchings*, Discrete Applied Mathematics **132** (2004) 163-174.
- [18] V. E. Levit, E. Mandrescu, *On α -critical edges in König-Egerváry graphs*, Discrete Mathematics **306** (2006) 1684-1693.

- [19] V. E. Levit, E. Mandrescu, *Triangle-free graphs with uniquely restricted maximum matchings and their corresponding greedoids*, Discrete Applied Mathematics **155** (2007) 2414–2425.
- [20] V. E. Levit, E. Mandrescu, *Some structural properties of very well-covered graphs*, Congressus Numerantium **186** (2007) 97–106.
- [21] V. E. Levit, E. Mandrescu, *Well-covered graphs and greedoids*, Proceedings of the 14th Computing: The Australasian Theory Symposium (CATS2008), Wollongong, NSW, Conferences in Research and Practice in Information Technology Volume **77** (2008) 89–94.
- [22] V. E. Levit, E. Mandrescu, *The clique corona operation and greedoids*, Combinatorial Optimization and Applications, Second International Conference, COCOA 2008, Lecture Notes in Computer Science **5165** (2008) 384–392.
- [23] V. E. Levit, E. Mandrescu, *Interval greedoids and local maximum stable sets in graphs*, (2008) arXiv:0811.4089v1[math.CO] 13 pp.
- [24] V. E. Levit, E. Mandrescu, *On local maximum stable sets of the corona of a path with complete graphs*, Proceedings of the 6th Congress of Romanian Mathematicians (2009) 565–569.
- [25] V. E. Levit, E. Mandrescu, *Graph operations that are good for greedoids*, Discrete Applied Mathematics **158** (2010) 1418–1423.
- [26] V. E. Levit, E. Mandrescu, *Very well-covered graphs of girth at least four and local maximum stable set greedoids*, Discrete Mathematics, Algorithms and Applications (2010) (accepted). arXiv:1008.2897v1[cs.DM] 7 pp.
- [27] G. L. Nemhauser, L. E. Trotter, Jr., *Vertex packings: structural properties and algorithms*, Mathematical Programming **8** (1975) 232–248.
- [28] M. D. Plummer, *Some covering concepts in graphs*, Journal of Combinatorial Theory **8** (1970) 91–98.
- [29] E. Prisner, J. Topp, P. D. Vestergaard, *Well-covered simplicial, chordal, and circular arc graphs*, Journal of Graph Theory **21** (1996) 113–119.
- [30] G. Ravindra, *Well-covered graphs*, J. Combin. Inform. System Sci. **2** (1977) 20–21.
- [31] J. A. Staples, *On some sub-classes of well-covered graphs*, Ph. D. Thesis, Vanderbilt University, 1975.
- [32] F. Sterboul, *A characterization of the graphs in which the transversal number equals the matching number*, Journal of Combinatorial Theory Series B **27** (1979) 228–229.
- [33] J. Topp, L. Volkmann, *On the well coveredness of products of graphs*, Ars Combinatoria **33** (1992) 199–215.